Author index

Ackerly, D.D. *see* Knight, C.A.

Allen, T.R., Millar, T., Berch, S.M. & Berbee, M.L. Culturing and direct DNA extraction find different fungi from the same ericoid mycorrhizal roots, 255

Andersen, C.P. *see* Lipp, C.C.

Anderson, L.J. *see* Withington, J.M.

Andrade, R.M. *see* Lambais, M.R.

Arnebrant, K. *see* Högberg, M.N.

Ashwath, N. *see* Bhatia, N.P.

Avis, P.G., McLaughlin, D.J., Dentinger, B.C. & Reich, P.B. Long-term increase in nitrogen supply alters above- and below-ground ectomycorrhizal communities and increases the dominance of *Russula* spp. in a temperate oak savanna, 239

Bååth, E. *see* Högberg, M.N.

Baker, A.J.M. *see* Bhatia, N.P.

Ballaré, C.L. *see* Robson, T.M.

Barlow, P.W. *see* Iijima, M.

BassiriRad, H. *see* Gutschick, V.P.

Batič, F. *see* Kranner, I.

Beckett, R.P. *see* Kranner, I.

Beever, R.E *see* Moyersoen, B.

Benech-Arnold, R.L., Enciso, S., Sánchez, R.A. & Rodríguez, M.V. On the hormonal nature of the stimulatory effect of high incubation temperatures on germination of dormant sorghum (*S.bicolor*) caryopses, 371

Bengough, A.G. *see* Iijima, M.

Berbee, M.L. *see* Allen, T.R.

Berch, S.M. *see* Allen, T.R.

Bhatia, N.P., Orlic, I., Siegele, R., Ashwath, N., Baker, A.J.M. & Walsh, K.B. Elemental mapping using PIXEshows the main pathway of nickel movement is principally symplastic within the fruit of the hyperaccumulator *Stackhousia tryonii*, 479

Bota, J. *see* Sampol, B.

Bouma, T.J. *see* Withington, J.M.

Brearley, F.Q., Press, M.C. & Scholes, J.D. Nutrients obtained from leaf litter can improve the growth of dipterocarp seedlings, 101

Bridge, P.D., Roberts, P.J., Spooner, B.M. & Panchal, G. On the unreliability of published DNA sequences, 43

Brown, M.T. *see* Nielsen, H.D.

Brownlee, C. *see* Nielsen, H.D.

Bułaj, B. *see* Withington, J.M.

Burke, T. *see* Jump, A.S.

Caldwell, M.M. *see* Robson, T.M.

Calfapietra, C., Gielen, B., Sabatti, M., De Angelis, P., Miglietta, F., Scarascia-Mugnozza, G. & Ceulemans, R. Do above-ground growth dynamics of poplar change with time under CO_2 enrichment?, 305

Carafa, A., Duckett, J.G. & Ligrone, R. Subterranean gametophytic axes in the primitive liverwort *Haplomitrium* harbour a unique type of endophytic association with aseptate fungi, 185

Ceulemans, R. *see* Calfapietra, C.

Cleyet-Marel, J.-C. *see* Larcher, M.

Coelho, S.M. *see* Nielsen, H.D.

Dancer, J. *see* Griffiths, R.G.

De Angelis, P. *see* Calfapietra, C.

DeLucia, E.H. *see* George, K.

Dentinger, B.C. *see* Avis, P.G.

Derner, J.D. *see* Polley, H.W.

Duckett, J.G. *see* Carafa, A.

Edmonds, R.L. *see* Trudell, S.A.

Eissenstat, D.M. *see* Withington, J.M.

Elkin, A.D. *see* Withington, J.M.

Enciso, S. *see* Benech-Arnold, R.L.

Flexas, J. *see* Sampol, B.

Flint, S.D. *see* Robson, T.M.

Flynn, K.J. *see* Stephens, N.

Gallon, J.R. *see* Stephens, N.

Gebauer, G. & Meyer, M. ^{15}N and ^{13}C natural abundance of autotrophic and myco-heterotrophic orchids provides insight into nitrogen and carbon gain from fungal association, 209

George, K., Norby, R.J., Hamilton, J.G. & DeLucia, E.H. Fine-root respiration in a loblolly pine and sweetgum forest growing in elevated CO_2, 511

Gielen, B. *see* Calfapietra, C.

Green, T.G.A. *see* Wirtz, N.

Griffiths, G. *see* Wagstaff, C.

Griffiths, R.G., Dancer, J., O'Neill, E. & Harwood, J.L. Lipid composition of *Botrytis cinerea* and inhibition of its radiolabelling by the fungicide iprodione, 199

Grimanelli, D. *see* Gurney, A.L.

Gurney, A.L., Grimanelli, D., Kanampiu, F., Hoisington, D., Scholes, J.D. & Press, M.C. Novel sources of resistanceto *Striga hermonthica* in *Tripsacum dactyloides*, a wild relative of maize, 557

Gutschick, V.P. & BassiriRad, H. Extreme events as shaping physiology, ecology, and evolution of plants: toward a unified definition and evaluation of their consequences, 21

Högberg, M.N., Bååth, E., Nordgren, A., Arnebrant, K. & Högberg, P. Contrasting effects of nitrogen availability on plant carbon supply to mycorrhizal fungi and saprotrophs – a hypothesis based on field observations in boreal forest, 225

Högberg, P. *see* Högberg, M.N.
Hamilton, J.G. *see* George, K.
Harwood, J.L. *see* Griffiths, R.G.
Hoisington, D. *see* Gurney, A.L.
Hu, C.-H. *see* Ryu, C.-M.
Huang, C.X. *see* Ryan, M.H.

Iijima, M., Barlow, P.W. & Bengough, A.G. Root cap structure and cell production rates of maize (*Zea mays*) roots in compacted sand, 127

Johnson, H.B. *see* Polley, H.W.
Joling, J. *see* Ritsema, T.
Jump, A.S. & Woodward, F.I. Seed production and population density decline approaching the range-edge of *Cirsium* species, 349
Jump, A.S., Woodward, F.I. & Burke, T. *Cirsium* species show disparity in patterns of genetic variation at their range-edge, despite similar patterns of reproduction and isolation, 359

Kõljalg, U. *see* Öpik, M.
Kanampiu, F. *see* Gurney, A.L.
Kloepper, J.W. *see* Ryu, C.-M.
Knight, C.A. & Ackerly, D.D. Evolution and plasticity of photosynthetic thermal tolerance, specific leaf area and leaf size: congeneric species from desert and coastal environments, 337
Kranner, I., Zorn, M., Turk, B., Wornik, S., Beckett, R.P. & Batič, F. Biochemical traits of lichens differing in relative desiccation tolerance, 167
Kyllo, D.A., Velez, V. & Tyree, M.T. Combined effects of arbuscular mycorrhizas and light on water uptake of the neotropical understory shrubs, *Piper* and *Psychotria*, 443

Lambais, M.R., Ríos-Ruiz, W.F. & Andrade, R.M. Antioxidant responses in bean (*Phaseolus vulgaris*) roots colonized by arbuscular mycorrhizal fungi, 421
Larcher, M., Muller, B., Mantelin, S., Rapior, S. & Cleyet-Marel, J.-C. Early modifications of *Brassica napus* root system architecture induced by a plant growth-promoting *Phyllobacterium* strain, 119
Leverentz, M. *see* Wagstaff, C.
Ligrone, R. *see* Carafa, A.
Liira, J. *see* Öpik, M.
Lipp, C.C. & Andersen, C.P. Role of carbohydrate supplyin white and brown root respiration of ponderosa pine, 523
Logan, D.C. Mitochondrial dynamics, 463
Lumbsch, H.T. *see* Wirtz, N.
Lusk, C.H., Wright, I. & Reich, P.B. Photosynthetic differences contribute to competitive advantage of evergreen angiosperm trees over evergreen conifers in productive habitats, 329
Lux, A. *see* Šottníková, A.

Malcolm, P. *see* Wagstaff, C.
Mantelin, S. *see* Larcher, M.
Martin, F. *see* Moyersoen, B.
Mattsson, M. & Schjoerring, J.K. Senescence-induced changes in apoplastic and bulk tissue ammonia concentrations of ryegrass leaves, 505
McCully, M.E. *see* Ryan, M.H.
McKinnon, L.M. & Mitchell, A.K. Photoprotection, not increased growth, characterizes the response of Engelmann spruce (*Picea engelmannii*) seedlings to high light, even when resources are plentiful, 69
McLaughlin, D.J. *see* Avis, P.G.
Medrano, H. *see* Sampol, B.
Meyer, M. *see* Gebauer, G.
Miglietta, F. *see* Calfapietra, C.
Millar, T. *see* Allen, T.R.
Mitchell, A.K. *see* McKinnon, L.M.
Modrzyński, J. *see* Withington, J.M.
Moora, M. *see* Öpik, M.
Moyersoen, B., Beever, R.E & Martin, F. Genetic diversity of *Pisolithus* in New Zealand indicates multiple long-distance dispersal from Australia, 569
Muller, B. *see* Larcher, M.

Nielsen, H.D., Brownlee, C., Coelho, S.M. & Brown, M.T. Inter-population differences in inherited copper tolerance involve photosynthetic adaptation and exclusion mechanisms in *Fucus serratus*, 157
Norby, R.J. *see* George, K.
Nordgren, A. *see* Högberg, M.N.
Novoplansky, A. Ecological implications of the determination of branch hierarchies, 111

O'Neill, E. *see* Griffiths, R.G.
Oleksyn, J. *see* Withington, J.M.
Olesiński, J. *see* Withington, J.M.
Öpik, M., Moora, M., Liira, J., Kõljalg, U., Zobel, M. & Sen, R. Divergent arbuscular mycorrhizal fungal communities colonize roots of *Pulsatilla* spp. in boreal Scots pine forest and grassland soils, 581
Orlic, I. *see* Bhatia, N.P.

Panchal, G. *see* Bridge, P.D.
Pancotto, V.A. *see* Robson, T.M.
Pearce, I.S.K., Woodin, S.J. & van der Wal, R. Physiological and growth responses of the montane bryophyte *Racomitrium lanuginosum* to atmospheric nitrogen deposition, 145
Pintado, A. *see* Wirtz, N.
Polley, H.W., Johnson, H.B. & Derner, J.D. Increasing CO_2 from subambient to superambient concentrations alters species composition and increases above-ground biomass in a C_3/C_4 grassland, 319
Press, M.C. *see* Brearley, F.Q.
Press, M.C. *see* Gurney, A.L.

Ríos-Ruiz, W.F. *see* Lambais, M.R.
Rafiq, A. *see* Wagstaff, C.

Rapior, S. *see* Larcher, M.
Read, J. & Sanson, G.D. Characterizing sclerophylly: the mechanical properties of a diverse range of leaf types, 81
Reddy, M.S. *see* Ryu, C.-M.
Reich, P.B. *see* Avis, P.G.
Reich, P.B. *see* Lusk, C.H.
Reich, P.B. *see* Withington, J.M.
Riera, D. *see* Sampol, B.
Ritsema, T., Joling, J. & Smeekens, S. Patterns of fructan synthesized by onion fructan : fructan 6G-fructosyltransferase expressed in tobacco BY2 cells – is fructan : fructan 1-fructosyltransferase needed in onion?, 61
Roberts, P.J. *see* Bridge, P.D.
Robson, T.M., Pancotto, V.A., Flint, S.D., Ballaré, C.L., Sala, O.E., Scopel, A.L. & Caldwell, M.M. Six years of solar UV-B manipulations affect growth of *Sphagnum* and vascular plants in a Tierra del Fuego peatland, 379
Rodríguez, M.V. *see* Benech-Arnold, R.L.
Rogers, H. *see* Wagstaff, C.
Ryan, M.H., McCully, M.E. & Huang, C.X. Location and quantification of phosphorus and other elements in fully hydrated, soil-grown arbuscular mycorrhizas: a cryo-analytical scanning electron microscopy study, 429
Rygiewicz, P.T. *see* Trudell, S.A.
Ryu, C.-M., Hu, C.-H., Reddy, M.S. & Kloepper, J.W. Different signaling pathways of induced resistance by rhizobacteria in *Arabidopsis thaliana* against two pathovars of *Pseudomonas syringae*, 413

Sánchez, R.A. *see* Benech-Arnold, R.L.
Sabatti, M. *see* Calfapietra, C.
Sala, O.E. *see* Robson, T.M.
Sampol, B., Bota, J., Riera, D., Medrano, H. & Flexas, J. Analysis of the virus-induced inhibition of photosynthesis in malmsey grapevines, 403
Sancho, L. *see* Wirtz, N.
Sanson, G.D. *see* Read, J.
Scarascia-Mugnozza, G. *see* Calfapietra, C.
Schjoerring, J.K. *see* Mattsson, M.
Scholes, J.D. *see* Brearley, F.Q.
Scholes, J.D. *see* Gurney, A.L.
Schroeter, B. *see* Wirtz, N.
Scopel, A.L. *see* Robson, T.M.
Sen, R. *see* Öpik, M.
Siegele, R. *see* Bhatia, N.P.
Smeekens, S. *see* Ritsema, T.
Šottníková, A. & Lux, A. Development, dilation and subdivision of cortical layers of gentian (*Gentiana asclepiadea*) root, 135
Spooner, B.M. *see* Bridge, P.D.
Stead, A. *see* Wagstaff, C.
Stephens, N., Flynn, K.J. & Gallon, J.R. Interrelationships between the pathways of inorganic nitrogen assimilation in the cyanobacterium *Gloeothece* can be described using a mathematical model, 545

Türk, R. *see* Wirtz, N.
Thomas, B. *see* Wagstaff, C.
Tracy, K.N. *see* Withington, J.M.
Trudell, S.A., Rygiewicz, P.T. & Edmonds, R.L. Nitrogen and carbon stable isotope abundances support the myco-heterotrophic nature and host-specificity of certain achlorophyllous plants, 391
Turk, B. *see* Kranner, I.
Tyree, M.T. *see* Kyllo, D.A.

Velez, V. *see* Kyllo, D.A.
Volaire, F. Seedling survival under drought differs between an annual (*Hordeum vulgare*) and a perennial grass (*Dactylis glomerata*), 501

Wagstaff, C., Malcolm, P., Rafiq, A., Leverentz, M., Griffiths, G., Thomas, B., Stead, A. & Rogers, H. Programmed cell death (PCD) processes begin extremely early in *Alstroemeria* petal senescence, 49
van der Wal, R. *see* Pearce, I.S.K.
Walsh, K.B. *see* Bhatia, N.P.
Webb, A.A.R. The physiology of circadian rhythms in plants, 281
Wirtz, N., Lumbsch, H.T., Green, T.G.A., Türk, R., Pintado, A., Sancho, L. & Schroeter, B. Lichen fungi have low cyanobiont selectivity in maritime Antarctica, 177
Withington, J.M., Elkin, A.D., Bułaj, B., Olesiński, J., Tracy, K.N., Bouma, T.J., Oleksyn, J., Anderson, L.J., Modrzyński, J., Reich, P.B. & Eissenstat, D.M. The impact of material used for minirhizotron tubes for root research, 533
Woodin, S.J. *see* Pearce, I.S.K.
Woodward, F.I. *see* Jump, A.S.
Wornik, S. *see* Kranner, I.
Wright, I. *see* Lusk, C.H.

Zobel, M. *see* Öpik, M.
Zorn, M. *see* Kranner, I.

Keyword Index

$\delta^{13}C$, 209
$\delta^{15}N$, 101, 209
^{13}C, 391
^{15}N, 391
A_L, 443
A_p, 443
abscisic acid, 371
acclimation, 4, 21
achlorophyllous plants, 391
aequorin, 281
algae, 157
Alstroemeria, 49
AM, 443
ammonia compensation point, 489
ammonium, 145, 489, 545
annual fine-root respiration, 511
apoplast, 489
Arabidopsis, 281
Arabidopsis thaliana, 413
arbuscular mycorrhiza, 443
arbuscular mycorrhizal (AM) fungi, 581
arbuscular mycorrhizal fungi (AMF), 429
arbuscular mycorrhizas, 185
Assimilation, 545

Banyalbufar malmsey (*Vitis vinifera*), 403
barley, 501
bioassay, 489
biogeographic limits, 21
biomass, 69
biomass production, 319
biotic stress, 403
border cell, 127
boreal forest, 225, 581
Borneo, 101
Botrytis cinerea, 199
branch competition, 111
branch hierarchy, 111
branch vigour, 111
British plants vegetation history, 280

C_3 species, 319
C_4 grasses, 319
calcium, 281, 429
Capronia, 255
carbon supply, 225
carotenoids, 69
Casparian band, 135
catalase (CAT), 421
cell production, 127
cell release, 127
Cenococcum, 239
chlorophyll, 69
chlorophyll fluorescence, 157
cholinephosphotransferase, 199
Circadian: stomata, 281
Cirsium, 349, 359
climate change, 21, 379
climate extremes, 4
clonal reproduction, 349, 359
CO_2 concentration, 319
colonization, 119
competition, 305, 329
conifers, 533
convergent adaptation, 337
Corallorhiza, 391
Cortinarius, 239
critical load, 145
cryo-scanning electron microscopy, 429
Cu^{2+}, 157
cyanobacteria, 177, 457, 545

Dactylis glomerata, 501
DAD-1 expression, 49
databases, 43
dehydration tolerance, 501
dehydrins, 501
denaturing gradient gel electrophoresis (DGGE), 581
Desiccation tolerance, 167
development, 279, 463
developmental decisions, 111
dicarboximide fungicide, 199
dilation growth, 135
dipterocarps, 101
dispersal, 569
disturbance response, 14
division, 463
DNA laddering, 49
DNA sequences, 5, 43, 462
drought resistance, 501
dynamics, 463

ectomycorrhiza, 391
ectomycorrhizal (ECM) fungi, 462
ectomycorrhizal fungi (EMF), 239
ectomycorrhizas, 101, 569
elemental mapping, 479
elevated CO_2, 7, 305
EMBL, 5
endangered plants, 581
endodermis, 135
environmental stress factors, 7
Ericaceae, 255
evolution, 21
exodermis, 135
extreme events, 21

fine root proliferation, 101
fine roots, 279, 533
fission, 463
flexure, 81
flora, 280
floristics, 280
fluorescence (F_v/F_m), 337
forest, 209
formononetin, 421
fracture, 81
free-air CO_2 enrichment (FACE), 305, 511
fructan, 61
fructan : fructan 6G-fructosyltransferase (6G-FFT), 61
fructosyltransferases, 61
Fucus serratus (Phaeophyta), 157
functional classes, 279
fungal communities, 239
fungal detection, 255
fungal diversity, 255
fungal ericoid mycorrhizal fungi, 255
fungi, 43
fusion, 463

GenBank, 5
genetic variation, 4, 359
gentian (*Gentiana asclepiadea*), 135
geographical range, 349, 359
geothermal, 569
germination, 371
gibberellins, 371
global change, 4
Gloeothece, 457
Gloeothece, 545
Glomeromycota, 185
Glomus clarum, 421
Glomus intraradices, 421
grapevine fan leaf virus (GFLV), 403
grapevine leaf roll-associated virus (GLRaV), 403
grapevines, 403
grassland, 209, 319

growth, 281, 305
growth rate, 329
growth respiration, 511
guaiacol-peroxidase (GPX), 421

Haplomitrium, 185
hardwoods, 533
haustorium, 557
heat shock protein, 337
heavy metal, 157
Hepaticae, 185
high pressure flow meter (HPFM), 443
Hordeum vulgare, 501
hormonal regulation, 371
host defence response, 429
hybridization, 17
hydraulic conductance, 443
Hymenoscyphus, 255
hyper-accumulation, 157

induced resistance, 413
introgression, 17
inulin neo-series, 61
iprodione, 199
isolation by distance, 359

K_t, 443
K_t/A_L: leaf specific conductance, 443
kestose, 61
Kunzea ericoides var. *microflora*, 569

latitudinal gradient, 349
leaf area (m^2), 443
leaf lifespan, 329
leaf litter, 101
leaf movements, 281
leaves, 14, 81
Leptospermum scoparium, 569
lichens, 177
light response, 523
lignin peroxidase, 462
lipids, 199
litter, 489
LMA, 14
Lobaria pulmonaria, 167
loblolly pine (*Pinus taeda*), 511
Lolium perenne (ryegrass), 489
long-term ecological study, 379
long-term experiments, 276

magnesium, 429
maintenance respiration, 511
maize, 461
maize (*Zea mays*), 127
manganese peroxidase, 462
maritime Antarctica, 177
measurements, 279
mesophyll limitations, 403
meta-analysis, 7
metal hyperaccumulation, 479
microbial biomass, 225
micro-PIXE, 479
microsatellites, 359
mid-cortex, 135
millet, 461
minirhizotrons, 533
mitochondria, 463
model, 545
modelling, 457
Monotropoideae, 391
montane heath, 145
morphology, 349, 463
myco-heterotrophic plants, 391
myco-heterotrophy, 209
mycorrhiza, 209
mycorrhizal fungi, 225

N_2 fixation, 457, 545
natural selection, 17, 21
neutrality, 21
New Zealand, 569
nickel (Ni), 479
nitrate reductase activity, 145
nitrate, 145, 545
nitrogen availability, 225
nitrogen concentration, 319
nitrogen deposition, 145
nitrogen fertilization, 239
nitrogen isotope discrimination, 101
nitrogen uptake respiration, 511
nitrogen use efficiency, 319
nitrogenase, 457
Nostoc, 177
N-status, 489
nuclear microprobe analysis, 479
nutrition, 69

oak savanna, 239
oilseed rape (*Brassica napus*), 119
onion (*Allium cepa*), 61
orchids, 209
Orobanche (broomrape), 461
osmotic adjustment, 501
ozone depletion, 276, 379

parasitic weeds, 461
parasitic plants, 557
PCR, 462
peatland, 379
Peltigera polydactyla, 167
peroxidase, 167
petal, 49
Phaseolus vulgaris (bean), 421
phenotypic plasticity, 111
phosphate, 421
phosphorus, 429
photoinhibition, 69
photosynthesis, 69, 157, 167, 329
photosynthetic metabolism, 403
photosynthetic rates, 111
phototransduction, 281
Phyllobacterium, 119
phylogenetic independent contrasts, 337
phylogenetics, 581
physiological ecology, 21
Picea engelmannii, 69
pigment, 167
Piper, 443
Pisolithus, 569
plant biomechanics, 81
plant community, 379
plant ecological strategies, 14
plant growth-promoting bacteria (PGPB), 119
plant growth-promoting rhizobacteria, 413
plant resistance, 557
plasticity, 337
polyploidy, 17
ponderosa pine (*Pinus poderosa*), 523
population density, 349
Populus, 305
potassium, 429
potassium leakage, 145
pre-harvest sprouting, 371
programmed cell death (PCD), 49
Pseudevernia furfuracea, 167
Pseudomonas syringae, 413
Psychotria, 443
public sequence databases, 5
Pulsatilla, 581

quantitative imaging, 479
quantitative trait, 349

Racomitrium lanuginosum, 145
rate-determined processes, 111
RDW, 443
realized environment, 4
redundant organs, 111
reliability, 43
reproductive responses, 7
rhizotron, 279

ribosomal RNA, 43
root cap, 127
root death, 279
root dry weight (g), 443
root cell vacuoles, 429
root excision, 523
root intercellular spaces, 429
root life span, 533
root meristem, 127
root morphology, 533
root pigmentation, 533
root respiration, 523
root survivorship, 533
root surface area (m^2), 443
root system architecture, 119
Russula, 239

saprotrophs, 225
sapwood density, 14
science policy, 7
sclerophylly, 81
Sebacina, 255
secondary thickening, 135
seed dormancy, 371
seed production, 349
seedling growth, 101
seedling, 581
seedling survival, 501
senescence, 49, 489
short rotation culture, 305
signalling, 281
soil compaction, 127
soil mechanical impedance, 127
soil pH, 225
soil sampling, 225
sorghum, 371, 461
speciation, 17
species composition, 319
specific leaf area, 329
specific leaf area (SLA), 337
specific root area ($m^2\ kg^{-1}$), 443
Sphagnum, 379
spot analysis, 479
SRA (A_f/RDW), 443
SSU rDNA sequences, 581
stable isotopes, 209, 391
stems, 14
stiffness, 81
stomata, 281
stomatal conductance, 329, 403
strength, 81
stress, 337
Striga hermonthica, 557
Striga (witchweed), 461
suberin lamella, 135
superoxide-dismutase (SOD), 421
sweetgum (*Liquidambar styeaciflua*), 511
symbiont specificity, 177
systematics, 43
taxonomy, 280
temperate forest, 511
temperature, 371
thermotolerance, 337
Tierra del Fuego, 379
tobacco BY2 cells, 61
tocopherol, 167
tolerance, 157
toughness, 81
Tripsacum dactyloides, 557
$tRNA^{Leu}$ intron, 177
tropical forest, 443
tropical rain forest, 101

ultrastructure, 49, 185
ultraviolet-B radiation, 276
UV-B, 276
UV-B radiation, 379

vegetation, 276
vegetation history, 280
voucher-based taxonomic studies, 5

wild relatives, 557

xanthophylls, 69
x-ray analysis, 479
X-ray microanalysis, 429